

BC Comparison of CIHI LTC Indicators with Canadian Average			
LTC Indicator	Measurement	BC average	Canada average
Daily Restraint Use	<u>Appropriateness and Effectiveness</u> Percentage of residents who were physically restrained daily as indicated on their target assessments (assessments used in the calculation)	11.6% (below average)	9.6%
Potentially Inappropriate Antipsychotic Use	<u>Appropriateness and Effectiveness</u> Percentage of residents on antipsychotics without a diagnosis of psychosis	33.2% (below average)	30.3%
Falls in the Last 30 Days	<u>Safety</u> Percentage of residents who fell in the 30 days prior to their target assessment (the assessment used in the calculation)	16.1% (below average)	14.9%
Worsened Pressure Ulcer	<u>Safety</u> Percentage of residents whose stage 2 to 4 pressure ulcer worsened since the prior assessment	3.2% (below average)	3.0%
Worsened Depressive Mood	<u>Health Status</u> Percentage of residents whose mood from symptoms of depression have worsened since the prior assessment	17.1% (above average)	23.8%
Improved Physical Functioning	<u>Health Status</u> The percentage of residents who improved or remained independent in transferring and locomotion (mid-loss ADLs) since the previous assessment	33.3% (above average)	32.2%
Worsened Physical Functioning	<u>Health Status</u> The percentage of residents who worsened or remained completely dependent in transferring and locomotion (mid-loss ADLs) since the prior assessment	31.3% (above average)	34%
Experiencing Pain	<u>Health Status</u> Percentage of residents who had pain as documented on the target assessment	16.7% (below average)	10.2%
Experiencing Worsened Pain	<u>Health Status</u> Percentage of residents whose pain worsened since prior assessment	10.6% (above average)	11.3%